Neuroscience Letters, 116 (1990) 393–397
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abe, T., see Toghi, H., 194
Aceves, J., see Floran, B., 136
Adamson, S., see Murphy, S., 7
Aizenman, E., Frost White, W., Loring, R.H. and Rosenberg, P.A., A 3,4-dihydroxyphenylalanine oxidation product is a non-*N*-methyl-D-aspartate glutamatergic agonist in rat cortical neurons, 168
Akasu, T., see Tokimasa, T., 304
Alterio, J., see Jacquemin, E., 23
Ando, S., see Waki, H., 367

Barnard, E.A., see Henley, J.M., 17
Berger, W., Horstmann, G.A. and Dietz, V., Interlimb coordination of stance in children: divergent modulation of spinal reflex responses and cerebral evoked potentials in terms of age, 118
Bernard, J.F., Villanueva, L., Carroué, J. and Le Bars, D., Efferent projections from the subnucleus reticularis dorsalis (SRD): a *Phaseolus vulgaris* leucoagglutinin study in the rat, 257
Beyreuther, K., see Dalaère, P., 87
Binns, K.E., see Whithington-Wray, D.J., 280
Böhmer, G., see Schmid, K., 204
Bohnett, M., see Messer, Jr., W.S., 184
Bohus, B., see Roozendaal, B., 210
Bremer, M.E., see Contreras, P.C., 190
Brooker, G., see Yakovlev, A.G., 216
Buhl, E.H. and Dann, J.F., Basal dendrites are a regular feature of hippocampal granule cells in flying fox hippocampus, 263
Burgard, E.C. and Sarvey, J.M., Muscarinic receptor activation facilitates the induction of long-term potentiation (LTP) in the rat dentate gyrus, 34

Cadet, J.L., see Przedborski, S., 51
Campbell, G., see Murphy, S., 7
Carroué, J., see Bernard, J.F., 257
Chen, J.C. and Chesler, M., A bicarbonate-dependent increase in extracellular pH mediated by GABAA receptors in turtle cerebellum, 130
Cheron, G., see Godaux, E., 162
Chesler, M., see Chen, J.C., 130
Chesselet, M.-F., see Litwak, J., 179
Chung, J.M., see Nam, S.C., 40
Chung, K., see Nam, S.C., 40
Chédotal, A. and Hamel, E., Serotonin-synthesizing nerve fibers in rat and cat cerebral arteries and arterioles: immunohistochemistry of tryptophan-5-hydroxylase, 269
Clough-Helfman, C., see Phillis, J.W., 315
Contreras, P.C., Bremer, M.E. and Gray, N.M., Ifenprodil and SL 82.0715 potently inhibit binding of [3H](+)-3-PPP to binding sites in rat brain, 190
Coireia, M.J. and Lang, D.C., A electrophysiological comparison of solitary type I and type II vestibular hair cells, 106
Corrette, B.J., see Skiebe, P., 261
Costa, E., see Yakovlev, A.G., 216
Courtois, Y., see Jacquemin, E., 23
Curry, K., see McLarnon, J.G., 341

Dann, J.F., see Buhl, E.H., 263
De Bernardi, M.A., see Yakovlev, A.G., 216
De Boer, P., Westerink, B.H.C. and Horn, A.S., The effect of acetylcholinesterase inhibition on the release of acetylcholine from the striatum in vivo: interaction with autoreceptor responses, 357
Delagrave, C., see Philippe, E., 12
Delaère, P., Duyckaerts, C., Masters, C., Beyreuther, K., Peitte, F. and Hauw, J.-J., Large amounts of neocortical A4 deposits without neuritic plaques nor tangles in a psychometrically assessed, non-demented person, 87
Denis, P., Elena, P.-P., Nordmann, J.-P., Saraux, H. and Lapalus, P., Autoradiographic localization of D1 and D2 dopamine binding sites in the human retina, 81
Deshpande, S.B. and Warnick, J.E., Interaction of thyrotropin-releasing hormone and methysergide on the monosynaptic reflex in isolated mammalian spinal cord, 141
Dietz, V., see Berger, W., 118
Dohi, T., see Kitayama, S., 275
Dudel, J., Depolarizing pulses to neuromuscular terminals of frogs can elicit graded, phasic transmitter release in the absence of Ca influx, 94
Duyckaerts, C., see Delaère, P., 87

Egner, U., see Harschuh, W., 245
Elena, P.-P., see Denis, P., 81
Emonet-Dénand, F., Laporte, Y. and Proske, U., Summation of tension in motor units of the soleus muscle of the cat, 112

0304-3940/90/$ 03.50 © 1990 Elsevier Scientific Publishers Ireland Ltd.

Fabrazzo, M., see Yakovlev, A.G., 216
Fahn, S., see Przedborski, S., 51
Fedele, E., see Raiteri, M., 352
Feng, D.-M., see Wiesenfeld-Hallin, Z., 293
Flament, D., see Hall, E.J., 379
Flock, A., see Karlsson, K.K., 101
Floran, B., Aceves, J., Sierra, A. and Martinez-Fong, D., Activation of D_1 dopamine receptors stimulates the release of GABA in the basal ganglia of the rat, 136
Folkers, K., see Wiesenfeld-Hallin, Z., 293
Fontana, G., see Raiteri, M., 352
Franck, J.E., see Newell, D.W., 325
Fraser, C., see Hall, E.J., 379
Frost White, W., see Aizenman, E., 168
Fujimoto, K., see Jibiki, I., 221

Gaulin, F., see Philippe, E., 12
Gaumann, D.M., Yaksh, T.L. and Lucas, D.L., Cryptic Met-enkephalin in adrenal and portal vein during splanchnic artery occlusion shock in cats, 387
Geffard, M., see Philippe, E., 12
Ghosh, A., see Mahata, S.K., 336
Godaux, E., Cheron, G. and Mettens, P., Ketamine induces failure of the oculomotor neural integrator in the cat, 162
Gramsbergen, J.B.P., see Speciale, C., 198
Gray, N.M., see Contreras, P.C., 190
Gross, R.A., see Kelly, K.M., 233
Guthrie, K.M., see Wilson, D.A., 250

Håkanson, R., see Wiesenfeld-Hallin, Z., 293
Hall, E.J., Flament, D., Fraser, C. and Lemon, R.N., Non-invasive brain stimulation reveals reorganised cortical outputs in amputees, 379
Halley, C., see Jacquemin, E., 23
Hallström, A., Sato, A., Sato, Y. and Ungerstedt, U., Effect of stimulation of the nucleus basalis of Meynert on blood flow and extracellular lactate in the cerebral cortex with special reference to the effect of noxious stimulation of skin and hypoxia, 227
Hamanaka, T., see Ota, M., 372
Hamel, E., see Chédotal, A., 269
Hartschuh, W., Weihe, E. and Egner, U., Electron microscopic immunogold cytochemistry reveals chromogranin A confined to secretory granules of porcine Merkel cells, 245
Hattori, Y., Moriwake, A. and Hori, Y., Biphasic effects of polarizing current on adenosine-sensitive generation of cyclic AMP in rat cerebral cortex, 320
Hauw, J.-J., see Delaère, P., 87
Hayashida, Y., see Kannan, H., 123
Heinemann, U., see Mudrick, L.A., 172
Henley, J.M. and Barnard, E.A., Autoradiographic distribution of binding sites for the non-NMDA receptor antagonist CNQX in chick brain, 17
Hennequin, S., see Morin, D., 299
Hilaire, G., see Morin, D., 299
Hori, Y., see Hattori, Y., 320
Horn, A.S., see De Boer, P., 357
Horstmann, G.A., see Berger, W., 118
Houtani, T., see Ueyama, T., 70
Howes, K.A., see Yu, H.-S., 309
Hughes, S. and Smith, M.E., Opioid receptors in skeletal muscle of normal and dystrophic mice, 29
Hutson, P.H., Semark, J.E. and Middlemiss, D.N., The TRH analogue MK-771, increases acetylcholine release in hippocampus but not striatum of the conscious rat, 149

Ikeda, M., see Ueyama, T., 70

Jacquemin, E., Halley, C., Alterio, J., Laurent, M., Courtois, Y. and Jeanny, J.C., Localization of acidic fibroblast growth factor (aFGF) mRNA in mouse and bovine retina by in situ hybridization, 23
Jeanny, J.C., see Jacquemin, E., 23
Jibiki, I., Fujimoto, K., Kubota, T. and Yamaguchi, N., MK-801 (non-competitive NMDA receptor antagonist)-induced blockade of the 'burst response' in the dentate gyrus after perforant path kindling in acutely prepared rabbits, 221
Joó, F., see Mihály, A., 45

Kannan, H., Ueta, Y., Nakamura, T., Yamashita, H. and Hayashida, Y., Effects of centrally administered atrial natriuretic polypeptide on sympathetic nerve activity and blood flow to the kidney in conscious rats, 123
Karlsson, K.K. and Flock, A., Quinine causes isolated outher hair cells to change length, 101
Keating, M.J., see Whithington-Wray, D.J., 280
Kelly, K.M., Gross, R.A. and Macdonald, R.L., Valproic acid selectively reduces the low-threshold (T) calcium current in rat nodose neurons, 233
Kim, K.J., see Nam, S.C., 40
Kitajiri, M., see Ueyama, T., 70

Kitayama, S., Ohtsuki, H., Morita, K., Dohi, T. and Tsujimoto, A., Bis-oxonol experiment on plasma membrane potentials of bovine adrenal chromaffin cells: depolarizing stimuli and their possible interaction, 275
Kon, K., see Waki, H., 367
Koolhaas, J.M., see Roozendaal, B., 210
Kozuki, K., see Ueyama, T., 70
Krug, M., see Mihály, A., 45
Kubota, T., see Jibiki, I., 221
Kuhnt, U., see Mihály, A., 45
Kumazawa, T., see Ueyama, T., 70

Lang, D.C., see Correia, M.J., 106
Lapalus, P., see Denis, P., 81
Laporte, Y., see Emonet-Dénand, F., 112
Laurent, M., see Jacquemin, E., 23
Le Bars, D., see Bernard, J.F., 257
Leem, J.W., see Nam, S.C., 40
Lemon, R.N., see Hall, E.J., 379
Leon, M., see Wilson, D.A., 250
Litwak, J., Mercugliano, M., Chesselet, M.-F., and Oltmans, G.A., Increased glutamic acid decarboxylase (GAD) mRNA and GAD activity in cerebellar Purkinje cells following lesion-induced increases in cell firing, 179
Loring, R.H., see Aizenman, E., 168
Lucas, D.L., see Gaumann, D.M., 387

Macdonald, R.L., see Kelly, K.M., 233
Mahata, S.K. and Ghosh, A., Neural influence on the action of insulin in the adrenomedullary catecholamine content in the pigeon, 336
Malouf, A.T., see Newell, D.W., 325
Marchi, M., Sanguineti, P. and Raiteri, M., Muscarinic receptors mediate direct inhibition of GABA release from rat striatal nerve terminals, 347
Martinez-Fong, D., see Floran, B., 136
Masters, C., see Delaère, P., 87
Matesz, C., Development of the oculomotor and trochlear nuclei in the *Xenopus* toad, 1
Matthies, H., see Mihály, A., 45
McLarnon, J.G. and Curry, K., Quinolinate activation of *N*-methyl-D-aspartate ion channels in rat hippocampal neurons, 341
Mercugliano, M., see Litwak, J., 179
Merkelbach, S., see Schmid, K., 204
Messer Jr., W.S., Bohnett, M. and Stibbe, J., Evidence for a preferential involvement of M1 muscarinic receptors in representational memory, 184
Mettens, P., see Godaux, E., 162
Middlemiss, D.N., see Hutson, P.H., 149
Mihály, A., Oláh, Z., Krug, M., Kuhnt, U., Matthies, H., Rapp, U.R. and Joó, F., Transient increase of *raf* protein kinase-like immunoreactivity in the rat dentate gyrus during long-term potentiation, 45
Mikkelsen, J.D., A neuronal projection from the lateral geniculate nucleus to the lateral hypothalamus of the rat demonstrated with *Phaseolus vulgaris* leucoagglutinin tracing, 58
Mocchetti, I., see Yakovlev, A.G., 216
Monteau, R., see Morin, D., 299
Morin, D., Hennequin, S., Monteau, R. and Hilaire, G., Depressant effect of raphe stimulation on inspiratory activity of the hypoglossal nerve: in vitro study in the newborn rat, 299
Morita, K., see Kitayama, S., 275
Moriwake, A., see Hattori, Y., 320
Morris, J.C., see Torack, R.M., 75
Mudrick, L.A. and Heinemann, U., Quisqualate-induced changes in extracellular sodium and calcium concentrations persist in the combined presence of NMDA and non-NMDA receptor antagonists in rat hippocampal slices, 172
Murphy, S., Osborne, P.B., Adamson, S. and Campbell, G., Co-localization of calcitonin gene-related peptide- and substance P-like immunoreactivity in mucosal intra-epithelial nerves in the toad colon, 7

Nagatsu, I., see Ueyama, T., 70
Nagatsu, T., see Ota, M., 372
Nakamura, T., see Kannan, H., 123
Nam, S.C., Kim, K.J., Leem, J.W., Chung, K. and Chung, J.M., Increased number of unmyelinated fibers in the ventral root after peripheral neurectomy in adult rat, 40
Naoi, M., see Ota, M., 372
Newell, D.W., Malouf, A.T. and Franck, J.E., Glutamate-mediated selective vulnerability to ischemia is present in organotypic cultures of hippocampus, 325
Nordmann, J.-P., see Denis, P., 81
Nozaki, Y., see Toghi, H., 194

Ohtsuki, H., see Kitayama, S., 275
Oldenburger, W.P., see Roozendaal, B., 210
Oltmans, G.A., see Litwak, J., 179
Oláh, Z., see Mihály, A., 45
Osborne, P.B., see Murphy, S., 7
Ota, M., Naoi, M., Hamanaka, T. and Nagatsu,

T., Inhibition of human brain aromatic L-amino acid decarboxylase by cooked food-derived 3-amino-1-methyl-5*H*-pyrido[4, 3-b]indole (Trp-P-2) and other heterocyclic amines, 372

Peitte, F., see Delaère, P., 87
Philippe, E., Gaulin, F., Delagrave, C. and Geffard, M., Expression of GABA-immunoreactivity by spinal motoneurons of some vertebrates, 12
Phillis, J.W. and Clough-Helfman, C., Protection from cerebral ischemic injury in gerbils with the spin trap agent *N*-tert-butyl-α-phenylnitrone (PBN), 315
Pittaluga, A., see Raiteri, M., 352
Pradet, A., see Verna, A., 156
Proske, U., see Emonet-Dénand, F., 112
Przedborski, S., Wright, M., Fahn, S. and Cadet, J.L., Quantitative autoradiographic changes in 5-[^{3}H]HT-labeled 5-HT_1 serotonin receptors in discrete regions of brain in the rat model of persistent dyskinesias induced by iminodipropionitrile (IDPN), 51

Raiteri, M., Fedele, E., Fontana, G. and Pittaluga, A., Glycine enhances [^{3}H]noradrenaline release from slices of rat hippocampus, 352
Raiteri, M., see Marchi, M., 347
Rapp, U.R., see Mihály, A., 45
Raso, R.S., Peripheral innervation of *Opsanus tao* (oyster toadfish), 241
Reiter, R.J., see Yu, H.-S., 309
Roeder, T., Histamine H1-receptor-like binding sites in the locust nervous tissue, 331
Roozendaal, B., Oldenburger, W.P., Strubbe, J.H., Koolhaas, J.M. and Bohus, B., The central amygdala is involved in the conditioned but not in the meal-induced cephalic insulin response in the rat, 210
Rosenberg, P.A., see Aizenman, E., 168
Roumy, M., see Verna, A., 156

Sanguineti, P., see Marchi, M., 347
Saraux, H., see Denis, P., 81
Sarvey, J.M., see Burgard, E.C., 34
Sato, A., see Hallström, A., 227
Sato, Y., see Hallström, A., 227
Schmid, K., Böhmer, G. and Merkelbach, S., Serotonergic control of phrenic motoneuronal activity at the level of the spinal cord of the rabbit, 204
Schwarcz, R., see Speciale, C., 198
Segade, L.A.G., Location of the motoneurons innervating the transverse mandibular muscle in the guinea pig, 64
Semark, J.E., see Hutson, P.H., 149
Shiraishi, M., see Tokimasa, T., 304
Sierra, A., see Floran, B., 136
Skiebe, P., Corrette, B.J. and Wiese, K., Evidence that histamine is the inhibitory transmitter of the auditory interneuron ON1 of crickets, 261
Smith, M.E., see Hughes, S., 29
Speciale, C., Wu, H.-Q., Gramsbergen, J.B.P., Turski, W.A., Ungerstedt, U. and Schwarcz, R., Determination of extracellular kynurenic acid in the striatum of unanesthetized rats: effect of aminooxyacetic acid, 198
Stibbe, J., see Messer Jr., W.S., 184
Strubbe, J.H., see Roozendaal, B., 210
Sugimoto, T., see Ueyama, T., 70
Sutor, B. and Ten Bruggencate, G., Ascorbic acid: a useful reductant to avoid oxidation fo catecholamines in electrophysiological experiments in vitro?, 287

Takahashi, J., see Toghi, H., 194
Takahashi, S., see Toghi, H., 194
Tanaka, Y., see Waki, H., 367
Ten Bruggencate, G., see Sutor, B., 287
Thalib, N., see Verna, A., 156
Toghi, H., Abe, T., Takahashi, S., Takahashi, J., Ueno, M. and Nozaki, Y., Effect of a synthetic norepinephrine precursor, L-threo-3,4-dihydroxyphenylserine on the total norepinephrine concentration in the cerebrospinal fluid of parkinsonian patients, 194
Tokimasa, T., Shiraishi, M. and Akasu, T., Morphological and electrophysiological properties of C-cells in bullfrog dorsal root ganglia, 304
Torack, R.M. and Morris, J.C., Tyrosine hydroxylase-like (TH) immunoreactivity in human mesolimbic system, 75
Tsujimoto, A., see Kitayama, S., 275
Turski, W.A., see Speciale, C., 198

Ueno, M., see Toghi, H., 194
Ueta, Y., see Kannan, H., 123
Ueyama, T., Kozuki, K., Houtani, T., Ikeda, M., Kitajiri, M., Yamashita, T., Kumazawa, T., Nagatsu, I. and Sugimoto, T., Immunolocalization of tyrosine hydroxylase and vasoactive intestinal polypeptide in nerve fibers innervating human palatine tonsil and paratonsillar glands, 70

Ungerstedt, U., see Hallström, A., 227
Ungerstedt, U., see Speciale, C., 198

Verna, A., Thalib, N., Roumy, M. and Pradet, A., Effects of metabolic inhibitors and hypoxia on the ATP and AMP content of the rabbit carotid body in vitro: the metabolic hypothesis in question, 156
Villanueva, L., see Bernard, J.F., 257

Waki, H., Kon, K., Tanaka, Y. and Ando, S., Age-related changes in the uptake of calcium channel blockers by brain capillary endothelial cells and synaptosomal fractions, 367
Warnick, J.E., see Deshpande, S.B., 141
Weihe, E., see Harschuh, W., 245
Wiesenfeld-Hallin, Z., Xu, X.-J., Håkanson, R., Feng, D.-M. and Folkers, K., Plasticity of the peptidergic mediation of spinal reflex facilitation after peripheral nerve section in the rat, 293
Westerink, B.H.C., see De Boer, P., 357
Whithington-Wray, D.J., Binns, K.E. and Keating, M.J., A four-day period of bimodality auditory and visual experience is sufficient to permit normal emergence of the map of auditory space in the guinea pig superior colliculus, 280
Wiese, K., see Skiebe, P., 261
Wilson, D.A., Guthrie, K.M. and Leon, M., Modification of olfactory bulb synaptic inhibition by early unilateral olfactory deprivation, 250
Wright, M., see Przedborski, W., 51
Wu, H.-Q., see Speciale, C., 198

Xu, X.-J., see Wiesenfeld-Hallin, Z., 293

Yakovlev, A.G., de Bernardi, M.A., Fabrazzo, M., Brooker, G., Costa, E. and Mocchetti, I., Regulation of nerve growth factor receptor mRNA content by dexamethasone: in vitro and in vivo studies, 216
Yaksh, T.L., see Gaumann, D.M., 387
Yamaguchi, N., see Jibiki, I., 221
Yamashita, H., see Kannan, H., 123
Yamashita, T., see Ueyama, T., 70
Yee, R.W., see Yu, H.-S., 309
Yu, H.-S., Yee, R.W., Howes, K.A. and Reiter, R.J., Diurnal rhythms of immunoreactive melatonin in the aqueous humor and serum of male pigmented rabbits, 309